Running Head: Physical activity and SCI

Developing Physical Activity Interventions for Adults with Spinal Cord Injury.

Part 2: Motivational Counseling and Peer-Mediated Interventions for

People Intending to be Active

BRIEF REPORT (265 LINES)

Amy E. Latimer-Cheung
Queen's University

Kelly P. Arbour-Nicitopoulos
University of Toronto

Lawrence R. Brawley
University of Saskatchewan

Casey Gray
CHEO Research Institute

A. Justine Wilson
University of British Columbia

Harry Prapavessis
Western University

Jennifer R. Tomasone
McMaster University

Dalton L. Wolfe
Lawson Health Research Institute

Kathleen A. Martin Ginis
McMaster University

Abstract

Objective: The majority of people with spinal cord injury (SCI) do not engage in sufficient leisure time physical activity (LTPA) to attain fitness benefits; however, many have good intentions to be active. This paper describes two pilot interventions targeting people with SCI who are insufficiently active but intend to be active (i.e., “intenders”).

Method: Study 1 examined the effects of a single, telephone-based counseling session on self-regulatory efficacy, intentions, and action plans for LTPA among 7 men and women with paraplegia or tetraplegia. Study 2 examined the effects of a home-based strength-training session, delivered by a peer and a fitness trainer, on strength-training task self-efficacy, intentions, action plans, and behavior. Participants were 11 men and women with paraplegia. **Results:** The counseling session (Study 1) yielded medium- to large-sized increases in participants’ confidence to set LTPA goals and intentions to be active. The home visit (Study 2) produced medium- to large-sized increases in task self-efficacy, barrier self-efficacy, intentions, action planning, and strength-training behavior from baseline to 4 weeks after the visit. **Conclusions/Implications:** Study 1 findings provide preliminary evidence that a single counseling session can impact key determinants of LTPA among intenders with SCI. Study 2 findings demonstrate the potential utility of a peer-mediated, home-based strength training session for positively influencing social cognitions and strength-training behavior. Together, these studies provide evidence and resources for intervention strategies to promote LTPA among intenders with SCI, a population for whom LTPA interventions and resources are scarcely available.

64 Keywords: exercise, social cognition, motivational interviewing, strength training, health
65 action process approach
66

Impact

- Study 1 is the first to demonstrate the efficacy of a brief motivational counseling session for strengthening social-cognitive determinants of physical activity in persons with SCI who intend to become active. Study 2 is the first study, in any population, to demonstrate the efficacy of employing peer mentors to deliver a brief intervention, in a person's home, to increase strength-training behavior and its social cognitive determinants.
- Given the dearth of research testing LTPA-enhancing interventions for persons with SCI, these studies provide preliminary evidence that both motivational counseling and peer-mediated strength-training sessions can enhance social cognitive determinants of LTPA. A peer-mediated session can also increase strength-training behavior.
- These pilot studies have resulted in the creation of evidence-based resources that can now be used by clinicians and service organizations to implement motivational counseling and peer-delivered LTPA-enhancing interventions.

1 The majority of people with spinal cord injury (SCI) are insufficiently active to
2 obtain fitness benefits (Martin Ginis, Latimer et al., 2010). However, as reported in the
3 first paper in this series (Martin Ginis, Tomasone, et al., 2011), many have good
4 intentions to become more active in the near future. This subset of individuals has been
5 labeled “intenders” (Schwarzer, 2008; Schwarzer, Lippke, & Luszczynska, 2011).
6 Intenders are an ideal target for leisure time physical activity (LTPA) interventions
7 because they are more ready for change than people who have no intention to change. In
8 this paper, we describe pilot testing of two interventions promoting LTPA in this group.

9 To date, only four published studies have evaluated the impact of LTPA-enhancing
10 interventions for people with SCI (Arbour-Nicitopoulos, Martin Ginis, & Latimer, 2009;
11 Latimer, Martin Ginis, & Arbour, 2006; Warm, Belza, Whitney, Mitchell, & Stiens,
12 2004; Zemper et al., 2003). These studies focused primarily on teaching self-regulation
13 strategies such as action planning and goal setting. In contrast, the interventions described
14 herein focus on enhancing theory-based determinants of LTPA—particularly self-
15 efficacy and intentions, as well as planning (Bandura, 1997; Schwarzer, 2008). Because
16 these social cognitions are known to influence LTPA in the SCI population (Arbour-
17 Nicitopoulos, Martin Ginis, Wilson, & The SHAPE-SCI Research Group, 2010; Latimer
18 & Martin Ginis, 2005; Martin Ginis, Latimer et al., 2011), and tend to be weaker among
19 intenders versus “actors” who are actively engaged in LTPA (Martin Ginis, Tomasone, et
20 al., 2011), they are optimal targets for interventions aimed at transitioning intenders into
21 actors (Schwarzer et al., 2011).

22 **Study 1: LTPA Motivational Counseling**

The first pilot study tested the efficacy of a single counseling session that applied motivational interviewing (MI) principles to strengthen social cognitions associated with LTPA. MI is a client-centered counseling approach that aims to enhance individuals' motivation and commitment to change (Miller & Rollnick, 2002). It typically consists of a few brief sessions where clients explore their values and motives for change (Hettema, Steele, & Miller, 2005). The counselor's role is to evoke the client's own motivation to change, rather than impose change. MI has been used to encourage LTPA participation in several clinical populations (Bennett, Lyons, Winters-Stone, Nail, & Scherer, 2007; Brodie & Inoue, 2005; Silva et al., 2008), but has not yet been utilized among people with SCI. Within the SCI community, MI has the potential to be an effective and efficient intervention technique in a number of contexts such as during a physician's visit or a telephone counseling session (Rubak, Sandbaek, Lauritzen, & Christensen, 2005). Using a single arm pre-test, post-test design with a one-day follow-up, we tested the hypothesis that a single counseling session that incorporates MI techniques would enhance intenders' confidence to schedule, to set goals, and to manage barriers for LTPA, as well as increase intentions and encourage implementation of an LTPA action plan.

Methods

Participants. Seven LTPA intenders were recruited from our participant database of people with SCI. Refer to Table 1 for study eligibility and Table 2 for participant demographic characteristics. Research Ethics Boards approved the protocol. All participants provided informed consent.

Measures. Demographic data were collected along with measures of the social cognitive variables (see Table 3). For descriptive purposes, minutes/week of moderate to

heavy intensity LTPA were assessed using the Leisure Time Physical Activity Questionnaire for People with SCI (LTPAQ-SCI; Martin Ginis & Latimer, 2007). (See Martin Ginis, Phang, Latimer, & Arbour-Nicitopoulos, 2011 for reliability and validity).

Intervention fidelity was assessed using the 6-item health-care climate questionnaire (Williams, Grow, Freedman, Ryan, & Deci, 1996). Items gauged the degree to which participants perceived the counselor as supportive versus controlling (e.g., "The physical activity coach provided me with choices and options"). Responses were made on a 7-point Likert-type scale. This tool has been validated to assess caregiver support for health-promoting behaviors such as LTPA, and shown acceptable reliability (Williams, Freedman, & Deci, 1998). Cronbach's alpha was 0.79 in the present study.

To assess intervention acceptability, participants indicated whether they found the session helpful (yes/no) and responded to a series of open-ended questions regarding specific intervention components (e.g., "what aspects were helpful/ unhelpful").

Procedure. All participant contact was via telephone. The project coordinator contacted potential participants to screen for eligibility. Eligible volunteers were scheduled to complete a counseling session. The interventionists were Exercise Psychology graduate students (1 MSc, 1PhD) who had completed a one-day motivational coaching course. Each 30-minute session began with activities to elicit participants' values, to build rapport between the participant and the interventionist, and to determine participants' willingness for change. The interventionist used strategies tailored to the clients' level of motivation to encourage them to resolve their uncertainty about change (see Table 3). She drew upon insights gained from the initial rapport-building activities and directed discussion toward topics of goal-setting, action planning, and scheduling

LTPA. She assisted participants in setting a concrete LTPA goal for the next week and discussed solutions to potential barriers to achieving this goal. One day later, the study coordinator administered the social cognitive and intervention fidelity measures and offered access to *Get In Motion*, a free physical activity telephone counseling service.

Results and discussion

Cohen's d effect sizes were calculated with effects of 0.20, 0.50, and 0.80 indicating small, medium, and large effects respectively (Cohen, 1992). Paired, one-tailed t -tests revealed significant medium- to large-sized increases in goal-setting self-efficacy, $d = 0.72$, and intention strength, $d = 1.01$, from pre- to post-intervention, $ps < .032$ (refer to Table 5). These positive findings are consistent with extant MI research in the general population (Knight, McGowan, Dickens, & Bundy, 2006; Martin & McNeil, 2009; Rubak et al., 2005), however many of those trials included multiple contacts lasting longer than 30 min each and incorporated supporting motivational materials. Our intervention used minimal contact and yet had significant positive effects. Demonstrating the impact of a minimal contact intervention has important implications in regard to potential integration of this approach in busy clinical settings.

Small- to medium-sized effects emerged for intentions, $d = 0.63$, and action planning, $d = 0.42$; however, these effects did not reach conventional levels of significance. There were small, non-significant decreases in scheduling self-efficacy, $d = -0.23$, and barrier self-efficacy, $d = -0.13$. The lack of statistical significance likely is the results of the small sample size and session content. It seems that the session enhanced participants' confidence in their ability to set realistic goals (goal-setting self-efficacy) and their commitment to these goals (intention strength) but it did not significantly

impact their willingness to increase the difficulty of their goals (intentions to increase physical activity participation). Additional session content related to goal progression may be beneficial. The lack of significant effects might also be indicative of the need for greater emphasis on action planning within the intervention. A greater focus on planning daily bouts of LTPA (action planning) and on planning for LTPA barriers (coping planning) may facilitate self-regulatory efficacy. For example, two randomized controlled trials evaluating the effectiveness of a telecounseling intervention for people with SCI that included directed discussion regarding LTPA action and coping planning, prevented decreases in confidence to scheduling LTPA and coping with barriers (Arbour-Nicitopoulos, et al., 2009; Latimer, et al., 2006).

Overall, the counseling session was well-received. The mean score on the healthcare climate questionnaire was 5.93 ± 0.72 out of 7, indicating that participants felt the counselor was supportive during the MI session. A tally of participants' open-ended responses revealed that 6/7 found the counseling session helpful. All participants indicated that the session was a good length of time. One participant liked having the session over the phone whereas two others would have preferred for the session to be in person. The remaining four participants did not indicate any preference.

Taken together, the results of our pilot test indicate that a single telephone-based session that includes MI techniques holds promise as a strategy for increasing intenders' confidence to set LTPA goals and strengthening their intentions to reach their goals.

Study 2: Peer-mediated home-based strength training intervention

To increase muscular strength, it is recommended that adults with SCI engage in strength-training exercises twice per week (Martin Ginis, Hicks et al., 2011). However,

less than 20% of people with SCI do any strength training whatsoever (Martin Ginis, Latimer, et al., 2010). Barriers to strength training include lack of accessible equipment in fitness centers (Arbour-Nicitopoulos & Martin Ginis, 2011), transportation barriers, and lack of strength-training exercise knowledge and confidence (Kasperavicius, et al., 2010). In the first study in this series (Martin Ginis, Tomasone, et al., 2011), we showed that intenders' confidence to perform strength-training activities was low. Accordingly, this study pilot tested an intervention designed to strategically address barriers and increase self-efficacy and strength-training behaviour among intenders with paraplegia.

The intervention entailed a single home visit by a certified personal trainer and a peer with paraplegia. Home visits are known to reduce barriers to accessing care and rehabilitation services (Cranen et al., 2011). Peers are a preferred source of LTPA information among people with SCI (Faulkner et al., 2009; Letts et al., 2011). Among able-bodied older adults, peer mentors have been shown to be effective for increasing LTPA through ongoing telephone counseling (Castro, Pruitt, Buman, & King, 2011). No published study to date—in any population—has examined the efficacy of employing peer mentors to deliver a brief, LTPA-enhancing intervention in a participant's home.

The intervention content, grounded in the self-efficacy aspect of Bandura's Social Cognitive Theory (1997), aimed to enhance task self-efficacy to perform strength-training activities by targeting the four sources of self-efficacy: mastery experience, vicarious experience, verbal persuasion and physiological feedback. Interventions targeting sources of self-efficacy, particularly mastery and vicarious experiences, have been shown to increase confidence for, and participation in, LTPA (for a review, see Ashford, Edmunds, & French, 2010; Williams & French, 2011). The intervention also aimed to increase

action planning by teaching this important skill, known to help people with SCI translate good intentions into LTPA (Arbour-Nicitopoulos, et al., 2009; Latimer, et al., 2006). Accordingly, we hypothesized that our peer-mediated, home-based strength-training intervention would increase self-efficacy, action planning, and strength-training behavior. This hypothesis was tested among intenders with paraplegia using a single arm pre-test, post-test design with a four-week follow-up.

Methods

Participants. Study eligibility criteria and participant characteristics are listed in Tables 1 and 2 respectively. Participants were recruited from a database, newsletter and website advertisements, and word-of-mouth. The Institutional Ethics Boards approved the protocol and all participants provided informed consent.

Measures. Refer to Table 3 for a description of the social-cognitive variables assessed. The LTPAQ-SCI was modified to measure only strength-training activity (as opposed to all LTPA). Participants recalled the frequency (# of bouts) and duration (min/bout) of moderate and heavy intensity strength-training activities over the past 7 days. The LTPAQ has demonstrated validity and reliability for measuring LTPA behavior including strength-training (Martin Ginis, Phang, et al., 2011).

Intervention fidelity was monitored using a visit checklist that the trainer completed following each home visit. The trainer recorded (yes/no) whether each of the prescribed intervention activities was completed and the duration of the home visit. Participants completed a 6-item scale assessing perceived benefits of the visit such as increased confidence to strength train (1= *definitely no*; 7= *definitely yes*).

Acceptability of the intervention was determined by assessing participants' satisfaction with the personal trainer (*Yes/No*), the peer (*Yes/No*), and the exercises (1= *very unsatisfied*; 7= *very satisfied*). Participants also rated (1= *definitely no*; 7= *definitely yes*) whether they could relate to the peer, found the peer helpful, and felt the home visit would help them meet their strength-training goals.

Procedure. The project coordinator screened participants via telephone. Those eligible were scheduled for a baseline interview, a home visit (1 wk later) and a follow-up interview (5 wks later). The coordinator administered the social cognitive and behavioral questionnaires by telephone at both interviews. Participants completed the intervention acceptability questionnaire immediately following the visit (online or paper copy).

Home visits were conducted by a personal trainer (Trainer 1: male, 5 visits; Trainer 2: female, 8 visits) and a peer with paraplegia (Peer 1: male, 5 visits; Peer 2: female, 7 visits, Peer 3: female, 1 visit). Trainers had 6-7 yrs experience training people with SCI. Peers were habitual exercisers drawn from a peer support program and an SCI exercise program. They were trained in intervention delivery by the first author (training manual available online); each trainer-peer dyad rehearsed before the home visit.

The activities included in the home visit are outlined in Table 5. The session began with an orientation that involved highlighting the benefits of strength training, discussing past mastery experiences, identifying existing resources for strength-training within the participant's home, and setting a strength-training goal. Next the peer modeled a set of seven exercises, the participant tried each exercise, and the trainer reinforced the participant's performance using verbally persuasive statements and by drawing upon examples of the participant's past mastery experiences. Finally, the trainer designed a

personalized strength-training program. The trainer and participant developed a one-week action plan for implementing the program. Participants received a copy of their personalized program, their action plan, a manual describing the exercises, and two different strengths of resistance bands.

Results and Discussion

Paired, one-tailed t-tests revealed significant medium to large-sized increases for task frequency self-efficacy, $d = 0.52$; barrier self-efficacy, $d = 0.87$; intentions, $d = 0.60$; and action planning, $d = 1.14$, $ps < .28$ (see Table 5). Although not significant, the intervention resulted in small to medium-sized increases in task duration self-efficacy, $d = 0.25$, and goal-setting self-efficacy, $d = 0.51$. Scheduling self-efficacy effects were negligible, $d = 0.07$. Significant large-sized effects emerged for each behavioral indicator. Number of bouts of strength-training, $d = 1.07$, bout duration, $d = 0.88$, and total min/week of strength-training, $d = 0.96$, increased significantly, $ps < .024$. At the four-week follow-up, 9 of 11 participants were strength training at least twice/week.

The magnitude of many of these positive effects is much larger than the magnitude of effects reported in two meta-analyses evaluating intervention techniques for changing self-efficacy and physical activity ($d = .03-.42$; Ashford et al., 2010; Williams & French, 2011). In these reviews, vicarious experience, providing instruction and feedback, and action planning were identified as being among the most effective strategies for enhancing self-efficacy and physical activity behavior. All of these techniques were featured prominently in our intervention and likely enhanced intervention impact.

The small effects for task duration, goal-setting, and scheduling self-efficacy likely are due to the timeline for follow-up and the intervention content. Follow-up was

conducted four weeks after the home visit. Increases in task duration self-efficacy may only emerge after several weeks of mastery experiences once participants experience noticeable training effects and develop muscular endurance. Furthermore, having the personal trainer determine the participants' goals may have undermined change in goal-setting self-efficacy. Although assigning goals can enhance task self-efficacy (e.g., Ashford et al., 2010; Elston & Martin, 2004), this may undermine participants' confidence to set and adjust goals on their own. Perhaps an additional session to foster this skill set is necessary. A subsequent session discussing strategies for fitting strength-training into a busy schedule also could be beneficial for enhancing scheduling self-efficacy (Arbour-Nicitopoulos et al., 2009).

Overall, participants evaluated the intervention favorably. Visits lasted 70.00 $\pm$ 19.52 min. Deviations from protocol related to shortening the exercise cool down (4 cases) and not distributing the end-of-visit survey (5 cases) because of time constraints. The 7 participants who completed the end-of-visit survey all strongly agreed (all M_s > 6.60 out of 7) they had gained confidence to perform the prescribed exercises, increased awareness of the benefits of strength training, knowledge and interest in home exercise options and learned new exercises and how to apply the information from the visit to their everyday routine. All were satisfied with the peer and personal trainer, and were very satisfied with the exercises ($M=6.50\pm0.54$ out of 7). Moreover, they felt strongly (M_s > 6.49 out of 7) that they could relate to the peer, that having the peer at the home visit was helpful, and that the visit would help them meet their strength training goals.

General Discussion

Our pilot studies provide evidence for two promising intervention strategies for encouraging LTPA participation among intenders with SCI. These studies also highlight a set of general principles to consider when designing interventions for intenders. First, according to HAPA, intenders have moved from the motivational to volitional phase of behavior change. However, it seems that there may be benefit in designing interventions to boost motivational determinants (e.g., self-efficacy) and motivation (i.e., intentions and intentions strength) in addition to enhancing their volition (i.e., action planning). Including both motivational and volitional intervention components in Study 2 led to large increases in strength-training behavior. As suggested from the trend for increased action planning following a motivation intervention in Study 1, and consistent with theory, boosting motivation should serve to enhance volition.

Second, the number of intervention sessions needed to promote change in social cognitions and behavior is another point for consideration. A single intervention contact can have an impact. Single sessions may be highly effective among intenders specifically because they are primed for change and a single session may be enough to “nudge” them into action. Although additional sessions would likely strengthen the effects (Knight et al., 2006; Rubak et al., 2005), there are practical settings where multiple intervention contacts are not feasible. One session may be adequate to elicit some change.

These recommendations must be considered in light of study limitations. Our analyses were underpowered; nonetheless, we demonstrated significant intervention effects. The sample sizes are comparable to those in other published LTPA intervention studies involving persons with SCI (e.g., $n = 12$; Warms et al., 2004). The lack of a control group prevents us from ruling out spontaneous (albeit unlikely) change in

outcomes, socially desirable responding, and non-systematic interventionist effects. A control group was not used because we felt it unethical to deny access to LTPA interventions in an underserved population where a standard care comparison group would entail no intervention whatsoever (Martin Ginis & Hicks, 2005). Measurement limitations included short follow-up periods, the absence of behavioral measures in Study 1, self-reported LTPA in Study 2 (although direct measures of LTPA do not exist), and the use of measures with emerging evidence of validity and reliability. Finally, the utility and feasibility of our intervention must be examined among intenders with tetraplegia.

Despite these limitations, our studies show the feasibility and potential impact of two intervention strategies for promoting LTPA among intenders with SCI. Furthermore, the resources developed for these studies have now been translated into evidence-based manuals, videos and guides that are publicly available (www.sciactioncanada.ca). Given that LTPA information and services are among the most desired but least available resources for people with SCI (Boyd & Bardak, 2004), these study outcomes are key milestones in addressing the needs of a terribly underserved community.

References

- Arbour-Nicitopoulos, K. P., Martin Ginis, K. A., & Latimer, A. E. (2009). Planning, leisure-time physical activity, and coping self-efficacy in persons with spinal cord injury: a randomized controlled trial. *Archives of Physical Medicine and Rehabilitation*, 90(12), 2003-2011.
- Arbour-Nicitopoulos, K. P., Martin Ginis, K. A., & Wilson, P. M. (2010). Examining the individual and perceived neighborhood associations of leisure-time physical activity in persons with spinal cord injury. *Annals of Behavioral Medicine*, 39(2), 192-197.
- Arbour-Nicitopoulos, K. P., & Martin Ginis, K. A. (2011). Examining the accessibility of fitness and recreational facilities for persons living with mobility disabilities. *Adapted Physical Activity Quarterly*, 28, 1-15.
- Ashford, S., Edmunds, J., & French, D. P. (2010). What is the best way to change self-efficacy to promote lifestyle and recreational physical activity? A systematic review with meta-analysis. *British Journal of Health Psychology*, 15(Pt 2), 265-288.
- Bandura, A. (1997). *Self-Efficacy: The Exercise of Control*. New York, NY: Freeman.
- Bennett, J. A., Lyons, K. S., Winters-Stone, K., Nail, L. M., & Scherer, J. (2007). Motivational interviewing to increase physical activity in long-term cancer survivors: a randomized controlled trial. *Nursing Research*, 56(1), 18-27.
- Boyd, E. & Bardak, L. (2004). *Cross Canada Checkup. Interim report of the national consultations on SCI services in Canada--a qualitative overview*. Rick Hansen Man in Motion Foundation.
- Brodie, D. A., & Inoue, A. (2005). Motivational interviewing to promote physical activity for people with chronic heart failure. *Journal of Advanced Nursing*, 50(5), 518-527.
- Castro, C. M., Pruitt, L. A., Buman, M. P., & King, A. C. (2011). Physical activity program delivery by professionals versus volunteers: the TEAM randomized trial. *Health Psychol*, 30(3), 285-294.
- Cohen, J. (1992). A power primer. *Psychological Bulletin*, 112(1), 155-159.
- Cranen, K., Drossaert, C. H., Brinkman, E. S., Braakman-Jansen, A. L., Ijzerman, M. J., & Vollenbroek-Hutten, M. M. (2011). An exploration of chronic pain patients' perceptions of home telerehabilitation services. *Health Expectations*. doi: 10.1111/j.1369-7625.2011.00668.x

- Elston, T., & Martin Ginis, K. A. (2004). The effects of self-set versus assigned goals on exercisers' self-efficacy for an unfamiliar task. *Journal of Sport & Exercise Psychology*, 26, 500-504.
- Faulkner, G., Gorczynski, P., Arbour, K., Letts, L., Wolfe, D., & Martin Ginis, K. A. (2009). Messengers and methods of disseminating health information among individuals with spinal cord injury: A scoping review. In T.C. Berkovsky (Ed.), *Spinal cord injuries: Types, treatments, and prognosis* (pp. 349-374). Hauppauge, NY: Nova Science Publishers.
- Gray, C., Wilson, A. J., Latimer, A. E., & Martin Ginis, K. A. (2010). *Motivational Interviewing and Physical Activity Manual*. Hamilton, ON: McMaster University.
- Hettema, J., Steele, J., & Miller, W. R. (2005). Motivational interviewing. *Annual Review of Clinical Psychology*, 1, 91-111.
- Kasperavicius, M. L., Latimer, A. E., Hetz, S. P., McColl, M. A., & Smith, K. (2010). A preliminary evaluation of a community-based exercise program for people with mobility impairments. In L. H. V. van der Woude, F. Hoekstra & S. de Groot (Eds.), *Assistive Technology Research Series* (pp. 273-275). Amsterdam, Netherlands: IOS press.
- Knight, K. M., McGowan, L., Dickens, C., & Bundy, C. (2006). A systematic review of motivational interviewing in physical health care settings. *British Journal of Health Psychology*, 11(Pt 2), 319-332.
- Lai, D. T., Cahill, K., Qin, Y., & Tang, J. L. (2010). Motivational interviewing for smoking cessation. *Cochrane Database Systematic Review*(1), CD006936. d
- Latimer, A. E., & Martin Ginis, K. A. (2005). The theory of planned behavior in prediction of leisure time physical activity among individuals with spinal cord injury. *Rehabilitation Psychology*, 50, 389-396.
- Latimer, A. E., Martin Ginis, K. A., & Arbour, K. P. (2006). The efficacy of an implementation intention intervention for promoting physical activity among individuals with spinal cord injury: A randomized controlled trial. *Rehabilitation Psychology*, 51, 273-280.
- Latimer, A. E., Martin Ginis, K. A., & Craven, B. C. (2004). Psychosocial predictors of exercise intentions and behavior among individuals with spinal cord injury. *Adapted Physical Activity Quarterly*, 21, 71-85.
- Marcus, B. H., & Simkin, L. R. (1993). The stages of exercise behavior. *Journal of Sports Medicine and Physical Fitness*, 33, 6.

- Martin Ginis, K. A., Arbour-Nicitopoulos, K. P., Latimer, A. E., Buchholz, A. C., Bray, S. R., Craven, B. C., et al. (2010). Leisure time physical activity in a population-based sample of people with spinal cord injury part II: activity types, intensities, and durations. *Arch Phys Med Rehabil*, 91(5), 729-733.
- Martin Ginis, K. A., & Hicks, A. L. (2005). Exercise research issues in the spinal cord injured population. *Exercise and Sport Science Reviews*, 33(1), 49-53.
- Martin Ginis, K. A., Hicks, A. L., Latimer, A. E., Warburton, D. E., Bourne, C., Ditor, D. S., et al. (2011). The development of evidence-informed physical activity guidelines for adults with spinal cord injury. *Spinal Cord*. doi: sc201163 [pii] 10.1038/sc.2011.63
- Martin Ginis, K. A., & Latimer, A. E. (2007). The Leisure Time Physical Activity Questionnaire for People with Spinal Cord Injury. *Available from the authors*.
- Martin Ginis, K. A., Latimer, A. E., Arbour-Nicitopoulos, K. P., Bassett, R. L., Wolfe, D. L., & Hanna, S. E. (2011). Determinants of physical activity among people with spinal cord injury: A test of social cognitive theory. *Annals of Behavioral Medicine*.
- Martin Ginis, K. A., Latimer, A. E., Arbour-Nicitopoulos, K. P., Buchholz, A. C., Bray, S. R., Craven, B. C., et al. (2010). Leisure time physical activity in a population-based sample of people with spinal cord injury part I: demographic and injury-related correlates. *Archives of Physical Medicine and Rehabilitation*, 91(5), 722-728.
- Martin Ginis, K. A., Latimer, A. E., Hicks, A. L., & Craven, B. C. (2005). Development and evaluation of an activity measure for people with spinal cord injury. *Medicine and Science in Sports and Exercise*, 37(7), 1099-1111.
- Martin Ginis, K. A., Latimer, A. E., Sinden, A. R., & Ginis, S. (2011). SCI Action Canada: A Framework for Developing and Mobilizing Physical Activity Interventions for People with Spinal Cord Injury. In S. Belanger, A. Aiken (eds). *Shaping the future, Military and Veterans Health Research*.
- Martin Ginis, K. A., Phang, S. H., Latimer, A. E., & Arbour-Nicitopoulos, K. P. (2011). Reliability and validity tests of the physical activity recall assessment for people with spinal cord injury (PARA-SCI) short form. *Archives of Physical Medicine and Rehabilitation*. *Available from authors*.
- Martin Ginis, K. A., Tomasone, J. R., Latimer, A. E., Arbour-Nicitopoulos, K. P., Bassett, R. L., & Wolfe, D. M. (2011). Development of theory-based physical activity interventions for people with spinal cord injury. *Rehabilitation Psychology*.

- Martins, R. K., & McNeil, D. W. (2009). Review of Motivational Interviewing in promoting health behaviors. *Clinical Psychology Review*, 29(4), 283-293.
- Miller, W. R., & Rollnick, S. (2002). *Motivational Interviewing: Preparing People For Change*. New York: The Guilford Press.
- Nash, M. S., Jacobs, P. L., Mendez, A. J., & Goldberg, R. B. (2001). Circuit resistance training improves the atherogenic lipid profiles of persons with chronic paraplegia. *Journal of Spinal Cord Medicine*, 24(1), 2-9.
- Rubak, S., Sandbaek, A., Lauritzen, T., & Christensen, B. (2005). Motivational interviewing: a systematic review and meta-analysis. *British Journal of General Practice*, 55(513), 305-312.
- Schwarzer, R. (2008). Modeling health behavior change: How to predict and modify the adoption and maintenance of health behaviors. *Applied Psychology-an International Review*, 57(1), 1-29.
- Schwarzer, R., Lippke, S., & Luszczynska, A. (2011). Mechanisms of health behavior change in persons with chronic illness or disability: the Health Action Process Approach (HAPA). *Rehabilitation Psychology*, 56(3), 161-170.
- Silva, M. N., Markland, D., Minderico, C. S., Vieira, P. N., Castro, M. M., Coutinho, S. R., et al. (2008). A randomized controlled trial to evaluate self-determination theory for exercise adherence and weight control: rationale and intervention description. *BMC Public Health*, 8, 234.
- Warms, C. A., Belza, B. L., Whitney, J. D., Mitchell, P. H., & Stiens, S. A. (2004). Lifestyle physical activity for individuals with spinal cord injury: A pilot study. *American Journal of Health Promotion*, 18, 288-291.
- Williams, G. C., Freedman, Z. R., & Deci, E. L. (1998). Supporting autonomy to motivate patients with diabetes for glucose control. *Diabetes Care*, 21(10), 1644-1651.
- Williams, G. C., Grow, V. M., Freedman, Z. R., Ryan, R. M., & Deci, E. L. (1996). Motivational predictors of weight loss and weight-loss maintenance. *Journal of Personality and Social Psychology*, 70(1), 115-126.
- Williams, S. L., & French, D. P. (2011). What are the most effective intervention techniques for changing physical activity self-efficacy and physical activity behaviour--and are they the same? *Health Education Research*, 26(2), 308-322.
- Wise, J. B., & Hale, S. B. (1999). Strengthening and generalizing self-efficacy in a male with a spinal cord injury. *Therapeutic Recreation Journal*, 33(4), 333-341.

448 Zemper, E. D., Tate, D. G., Roller, S., Forchheimer, M., Chiodo, A., Nelson, V. S., et al.
449 (2003). Assessment of a holistic wellness program for persons with spinal cord
450 injury. *American Journal of Physical Medicine and Rehabilitation*, 82(12), 957-
451 968.

452
453

454

Table 1

Inclusion Criteria

Eligibility Criteria	Study 1	Study 2
Age	18-65 years	18-65 years
Years Post-Injury	at least 12 months post	at least 6 months post
Injury Level	SCI resulting in tetraplegia or paraplegia	SCI resulting in paraplegia only ^a
Cognitive Impairment	None	None
Language Spoken	English	English
Location of Residence	Ontario, Canada	Within a 100 km radius of two Ontario cities ^b
Intender Status	Not engaging in regular moderate to heavy intensity LTPA (3 days/wk) as determined using the LTPAQ-SCI and had some intention to engage in LTPA over the next two weeks (i.e., scored >4 on the intentions to engage in LTPA once a week).	^c Not engaging in a regular (30 min 3 days/wk) strength-training program but intend to over the next 6 months as determined using the staging questionnaire (Marcus & Simkin, 1993).
Medical Screening	--	Free of medical concerns that prohibit participation in LTPA (i.e., self-reported doctor approval for participating in LTPA)

Note. ^aOnly participants with paraplegia were recruited for Study 2. The focus on people with paraplegia allowed us to evaluate the feasibility of the intervention approach while controlling for additional barriers related to limited hand function and the consequent need for additional assistance among some people with tetraplegia. ^bGeographical limits were imposed to reduce travel for our intervention teams. ^cThe criterion of strength training 3 days/wk was based on previous research (Latimer, et al., 2006) and was set prior to the release of SCI-specific physical activity guidelines advocating strength-training at least twice a week (Martin Ginis, Hicks, et al., 2011).

Table 2

Participant demographic characteristics

	Study 1 (<i>n</i> = 7)	Study 2 (<i>n</i> = 12)
Sex (% male)	57.14% (<i>n</i> = 4)	41.67% (<i>n</i> = 5)*
Age	51.86 ± 8.55 yrs	42.92 ± 15.61 yrs
Years post-injury	28.76 ± 25.35 yrs	23.31 ± 18.50 yrs
Injury level (% paraplegia)	85.71% (<i>n</i> = 6)	100% (<i>n</i> = 12)
Injury severity (% complete)	57.14% (<i>n</i> = 4)	58.33% (<i>n</i> = 7)
Mode of mobility (% manual chair users)	71.42% (<i>n</i> = 5)	75.00% (<i>n</i> = 9)
Education (% completed some post-secondary)	57.14% (<i>n</i> = 4)	100% (<i>n</i> = 12)

Note. *Two men could not be reached for follow-up in Study 2.

478 Table 3

479 *Social cognitive variables measured in Study 1 and Study 2.*

Variable	Brief description of items	Alpha	Reference
Intentions			
Study 1	3-items assessing intentions to engage in ≥ 15 mins of moderate to heavy intensity LTPA once, twice, three times per week over the next two weeks (1= <i>strongly disagree</i> , 7= <i>strongly agree</i>).	.76-.92	(Arbour-Nicitopoulos, et al., 2009)
Study 2	Same as Study 1, although items related to participating in ≥ 30 mins of strength-training.	.83-.95	
Intention strength			
Study 1	3-items where participants indicated the strength of their intentions to engage in ≥ 15 mins of moderate to heavy intensity LTPA once, twice, three times per week over the next two weeks (1 = <i>definitely will not</i> / 9 = <i>definitely will</i>)		
Action Planning			
Study 1	4-items evaluating whether participants had a detailed plan regarding when, where, how, and how often to engage in moderate to heavy intensity LTPA over the next two weeks (1= <i>completely disagree</i> ; 4= <i>completely agree</i>).	.60-.90	(Arbour-Nicitopoulos, et al., 2009)
Study 2	4-items evaluating whether participants had a detailed plan regarding when, what, where, and how to engage in strength training over the next 2 weeks (1= <i>definitely false</i> ; 7= <i>definitely true</i>).	.83-.99	
Goal-Setting SE			
Study 1	3-items assessing confidence over the next 2 weeks to: a) set realistic goals for engaging in ≥ 15 mins of moderate to heavy intensity LTPA, b) set realistic goals for maintaining a regular LTPA routine of ≥ 15 mins of moderate to hard intensity LTPA, and c) develop a plan for reaching LTPA goals (1= <i>not confident at all</i> ; 7= <i>completely confident</i>).	.78-.88	(Martin Ginis, Latimer, Arbour-Nicitopoulos, et al., 2011)
Study 2	Same as Study 1 with one additional item (4-items in total) assessing confidence to follow through with strength-training goals even though it may be difficult at times. The goal behavior was strength training 3x/wk for 30 mins each time.	.73-.94	

480

Scheduling			
SE	1-item assessing participants' confidence to manage	--	
Study 1	their weekly schedule to include ≥ 15 mins of moderate to heavy intensity LTPA over the next 2 weeks" (1= <i>not confident at all</i> ; 7= <i>completely confident</i>).		(Martin Ginis, Latimer, Arbour-
		.75-	Nicitopoulos,
Study 2	4-items assessing participants confidence to fit 30 mins of moderate to heavy intensity strength training into their weekly schedule once, twice, three times, and > three times per week over the next 2 weeks (1= <i>not confident at all</i> ; 7= <i>completely confident</i>).	.91	et al., 2011)
Barrier SE			
Study 1	9-items evaluating participants' confidence to overcome nine barriers commonly faced by individuals with SCI such as fatigue, time constraints, and pain (1= <i>not confident at all</i> ; 7= <i>completely confident</i>).	.93-.94	(Latimer, et al., 2006)
Study 2	5-items evaluating participants' confidence to overcome barriers to home-based strength training including: performing strength training exercises correctly, performing the exercises safely, adapting activities, designing a program, and coordinating resources		(Wise & Hale, 1999)
Task SE			
Study 2	<i>Duration</i> : Confidence to physically do moderate- and heavy-intensity strength training for 10-, 20-, 30-, 45-, and 60-mins, without stopping.	.82 to .95	(Martin Ginis, Latimer, Arbour-
	<i>Frequency</i> : Confidence to physically do 30 mins of moderate to heavy intensity strength training: once, twice, three times, > three times per week (1= <i>not confident at all</i> ; 7= <i>completely confident</i>).		Nicitopoulos, et al., 2011)

481

482

Table 4.

Motivational interviewing techniques used to promote change

Motivation to change	Counseling Strategies
Resistant to change	<ul style="list-style-type: none"> • Ask/discuss what is LTPA • Ask/discuss the benefits of LTPA • Ask/discuss the benefits of LTPA for people with SCI
Unsure about Change (ambivalent)	<ul style="list-style-type: none"> • Explore the status quo vs. change • Ask about what they know about LTPA • Ask about what activities they enjoy/don't enjoy doing • Ask about any particular health concerns they may have
Ready for Change	<ul style="list-style-type: none"> • Assist in choosing an activity- ask what they want to do before providing information • Assist in goal-setting • Address barriers and create coping plans for common barriers • Elicit confidence for change (seek support, help develop self-efficacy) • Discuss how to incorporate LTPA safely

487 Table 5.

488 *Social cognitive variables measured in Study 1 and Study 2.*

	Study 1				Study 2			
	Pre	Post	<i>d</i>	One-tailed t-test	Pre	Post	<i>d</i>	One-tailed t-test
Goal-Setting SE	5.69 ±0.93	6.24 ±0.54	0.72	$t(6) = -2.30, p = .031$	5.05 ±1.34	5.68 ±1.15	0.51	$t(10) = -1.46, p = .087$
Scheduling SE	6.14 ±0.69	5.86 ±1.21	-0.23	$t(6) = 0.80, p = .25$	5.18 ±1.68	5.27 ±1.01	0.07	$t(10) = -.18, p = .86$
Barrier SE	5.17 ±1.76	4.87 ±1.66	-0.13	$t(6) = 1.25, p = .44$	5.04 ±1.34	5.68 ±1.15	0.87	$t(10) = -2.17, p = .027$
Task SE								
Frequency					4.85 ±1.68	5.60 ±1.15	0.52	$t(9) = -2.32, p = .023$
Duration					4.78 ±2.01	5.32 ±2.30	0.25	$t(10) = -0.71, p = .25$
Intentions	5.57 ±1.70	6.38 ±0.61	0.63	$t(6) = -1.43, p = .10$	5.10 ±1.81	6.03 ±1.24	0.60	$t(10) = -3.10, p = .007$
Intentions strength ^a	6.62 ±1.99	8.24 ±1.08	1.01	$t(6) = -2.84, p = .015$				
Action planning ^b	2.89 ±0.99	3.23 ±0.58	0.42	$t(6) = -0.67, p = .26$	2.84 ±1.77	5.14 ±2.23	1.14	$t(10) = -3.55, p = .003$
Strength training								
Frequency (bouts/wk)					1.09 ±1.30	2.91 ±2.02	1.07	$t(10) = -2.71, p = .011$
Duration (min)					15.45 ±20.79	40.68 ±34.89	0.88	$t(10) = -2.30, p = .023$
Volume (min/wk)					30.45 ±37.98	82.50 ±66.70	0.96	$t(10) = -2.66, p = .012$

489 *Note.* ^aIntentions strength in Study 1 ranged from 1-9; ^bAction planning ranged from 1-5 in Study 1 and 1-7 in Study 2.

Table 6.

Intervention activities included in Study 2

Intervention component	Description	Theoretical basis
Orientation component		
Discuss activity preferences	Complete the Activity Preference and Motivation Worksheet drawing upon participants' previous experience with activities of daily living (ADL) with movement patterns similar to strength training.	Mastery experience
Discuss goals	Trainer set a goal for the participant to work up to performing strength training three times per week. The trainer assured participants that they could reach the goal given their current ability to perform a variety of ADL with similar movement patterns. Participants signed a goal commitment contract.	Verbal persuasion, mastery experience, goal setting
Environmental Assessment	The trainer, peer, and participant toured the participants' home to identify materials and locations that could be used for strength-training	Verbal persuasion
Strength-training component		
	The trainer and the peer guided the participant through short warm up and a series of seven strength-training exercises. The trainer introduced each exercise relating it to an ADL, had the peer demonstrate the exercises, and then encouraged the participant to perform 10 repetitions of the exercise. Participants rated the intensity of the exercise to provide the training feedback for designing the personalized program. Finally, the peer and the participant performed a cool-down.	Mastery experience, vicarious experience, verbal persuasion, physiological feedback
Planning component		
	The trainer provided the participant a written copy of a personalized strength-training program. The trainer and the participant developed a plan for implementing the program and discussed how to increase safely the intensity of the strength-training activities.	Verbal persuasion, goal setting